

## BIBLIOGRAPHY

Bjartell A, Abrahamsson PA, Björk T, Di Sant'agnese A, Matikainen MT, Lilja H. Production of alpha1-antichymotrypsin by PSA-containing cells of human prostate epithelium. *Urol.* 1993;42/5:502-10.

Björk T, Hulkko S, Bjartell A, Di Sant'agnese A, Abrahamsson PA, Lilja H. Alpha1-antichymotrypsin production in PSA-producing cells is common in prostate cancer but rare in benign prostatic hyperplasia. *Urol.* 1994;43/4:427-34.

Christensson A, Björk T, Nilsson O, et al. Serum Prostate Specific Antigen complexed to  $\alpha$ 1-antichymotrypsin as an indicator of prostate cancer. *J Urol.* 1993;150:100-5.

Lilja H. Significance of different molecular forms of serum PSA. *Urol Clin North Am.* 1993;20/4:681-6.

Lilja H, Christensson A, Dahlén U, et al. Prostate-Specific Antigen in serum occurs predominantly in complex with  $\alpha$ 1-antichymotrypsin. *Clin Chem.* 1991;37/9:1618-25.

Monda JM, Barry MJ, Oesterling JE. Prostate Specific Antigen cannot distinguish stage T1a (A1) prostate cancer from benign prostatic hyperplasia. *J Urol.* 1994;151:1291-5.

Sershon PD, Barry MJ, Oesterling JE. Serum Prostate-Specific Antigen discriminates weakly between men with benign prostatic hyperplasia and patients with organ-confined prostate cancer. *Eur Urol.* 1994;25:281-7.

Wu JT. Assay for Prostate Specific Antigen (PSA) : problems and possible solutions. *J Clin Lab Analysis.* 1994;8:51-62.